

# Understanding the Physical Processes

## Underlying Solar Eruptions

S. Antiochos, C.R. DeVore, J. T. Karpen, S. Masson, S. Guidoni (GSFC),  
T. Gombosi, B. van der Holst, W. D. Manchester, I. Sokolov, E. Landi, R. Frazin (U Mich)

- Partnership between GSFC and U Michigan to understand the most important drivers of space weather

### Approach:


- Develop MSEC (Modular Solar Eruptions Capability)
  - Second-generation NASA/LWS Strategic Capability designed for exploratory science by the general community
  - Designed for easy user modification and for local production runs (source code available)
- Build on heritage of SWMF
  - GB, *global background* module in which eruptions propagate
  - AR, high-res *active region* module in which eruptions originate
  - EE, *eruptive event* generator module
  - OB, *observables library* for generating SDO, STEREO, ACE, ... measurements from simulations
  - TR, *training library* for new users

## Science Objective:

- Flares & CMEs, (primary drivers of space weather)

## SDO observations of 08/31/12 eruptive flare:

- Pre-eruption structure always contains long sheared filament channel
- Balance between upward B pressure and downward tension
- System destabilizes, fast rise coincident with flare brightening
- Magnetic energy released into mass motions, heating, and energetic particles
- Filament and overlying corona ejected into solar wind



# Understanding Solar Eruptions


- How does the energy build up?
  - Why does magnetic shear concentrate at polarity inversion lines
  - Theories: flux emergence, flux cancellation, helicity condensation
- What destabilizes the system?
  - Ideal instability: kink or torus or loss of equilibrium
  - Resistive instability (reconnection): breakout or tether-cutting
- What causes the explosive energy release?
  - Ideal instability/loss of equilibrium or fast flare reconnection
- Where does the energy go?
  - Heating, mass acceleration, energetic particles, ...
  - Need accurate models for reconnection (*Toth*)
- What are the space weather impacts?
  - Need most accurate possible models of development and heliospheric propagation (*van der Holst, Manchester*)

# How does the energy build up?


## Flux Emergence:

- Often observe filament channel formation as part of emergence process (e.g., Okamoto et al 2009, Lites et al 2010)
- Emergence of subsurface twisted flux rope leads to sheared arcade at PIL (e.g., Manchester 2001, 2003, ...; Fan 2001, Magara et al 2003, ..)
- Continued emergence/cancellation increases shear (Manchester et al 2004, Fang et al 2012)
- In principle, may lead to eruption, but need larger simulations

(Simulation from Fang et al 2013)



Hinode SOT observation 12/06







# How does the energy build up?

## Helicity Condensation: (Antiochos 2013)

- Helicity injected into closed corona must be conserved
- Net helicity injected due to hemispheric rule
- Closed loops with same sense of twist merge by reconnection
- Opposite sense “bounce” (Linton & Antiochos)
- **Helicity** (shear) **condenses** onto PIL and CH boundary
- Produces filament channel and slow wind



# How does the energy build up?

- Rate of shear buildup at PIL given by:

$v_L = v_p (d / L_I)$ , where  $L_I = \sqrt{L^2 - H^2}$ ,  $d$  is scale of motions,  $v_p$  is their speed,  $L$  is scale of PIL, and  $H$  scale of CH

- Helicity spectrum:

$\theta_\lambda \sim \lambda^{-2}$ , except at largest scale  $L$  where shear builds up

- Physical origin of shear buildup at CH boundary:

$$H_T = \Phi_S \Phi_{L1} + \Phi_S \Phi_{CH}$$



But coronal hole cannot contribute to helicity; therefore, need canceling helicity:

$$H_{CH} = - \Phi_S \Phi_{CH}$$

# Modeling Helicity-Condensation





(Zhao et al., Knizhnik et al)

- Start with usual uniform field between two photospheric plates, a la Parker
- Apply slow twists at plates  $\sim$  supergranular flows
- Specify various cases for number and extent of flows
- “PIL” and “CH” given by boundaries of flow region
- Use *ARMS* MHD code to calculate evolution
- Helicity conserved to excellent accuracy



# Modeling Helicity-Condensation





Interaction of two  
twisted coronal  
loops:  
For same twist see  
reconnection and  
merger, opposite  
twist only kink  
(from Zhao, DeVore  
and Antiochos, 2014)





# Modeling Helicity-Condensation


Interaction of 84 twisted coronal loops (Knizhnik, DeVore & Antiochos 2014)



# Filament Channel Formation

(Mackay, DeVore, & Antiochos 2013)

- Magneto-frictional model for global field evolution
  - Includes diff. rotation, meridional flow, surface diffusion (Van Ballegooijen & Mackay)
- Wrong chirality for E-W PIL
- Added supergranular helicity injection and condensation to model
- Yields correct hemispheric rule for sufficiently large helicity condensation
- Also yields sheared filament channel rather than highly twisted flux rope




# What destabilizes system?

## Need explosive removal of overlying tension

(Sturrock, 2002)

- **Ideal instability/LOE:** (e.g., [Forbes](#) et al, Low, van Ballegooijen et al, Sturrock, Mikic et al, Roussev et al, Fan et al, ...)

- Filament is topologically distinct, twisted flux rope prior to eruption




- **Resistive:** (e.g., Moore et al, Antiochos et al, Aulanier, MacNeice et al, Lynch, ...)

- Filament is topologically part of overlying system

- Reconnection changes topology, removing overlying field

Karpen et  
al 2012



We (the community) will explore all mechanisms!

# Build up and destabilization

- Both flux emergence and helicity condensation can produce the free energy required for CME/flares
- Will model emergence for AR scale system and verify eruption
  - Determine whether ideal or reconnection-driven
  - Determine predictors for eruption
  - Test/validate with Hinode & SDO vector B observations
- Will model helicity condensation onto PILs and coronal holes
  - Determine whether results in MHD eruption
  - Test/validate with SDO vector B observations
  - Determine predictions for SPP & SO




# What Causes Explosive Energy Release?

## Fast Flare Reconnection


(Guidoni et al 2014)

- Evolution slow and weakly energetic until flare reconnection onset
- Flare reconnection releases bulk of energy: CME acceleration, particle acceleration, heating, shock
- Dynamics dominated by island formation
- Kinetic processes critically important




- Extension to fully 3D system
- Include solar wind
- Breakout reconnection has major implications for particle escape
- Explains impulsive SEP

(Masson et al 2014)




# What Causes Explosive Energy Release? **M**

- In 3D have multiple reconnections with neighboring flux systems
- Produces coronal dimmings
- “End” effects due to finite length for eruption




- Reconnection between erupting flux rope and neighboring open flux
- Allows flare particles to escape
- For SEP prediction, need accurate models for global coronal topology






# Energy Evolution





- Time history of magnetic and kinetic energies
  - CME onset corresponds to start of breakout reconnection
  - “Take-off” corresponds to start of fast flare reconnection
- Explosiveness all due to flare reconnection

# Eruption Mechanism



- Magnetic energy vs energy of corresponding “open” state
  - No evidence for ideal instability/loss of equilibrium
  - Resistive instability

# Where does the energy go?



- Basic onset and explosive evolution unchanged with  $S$
- Eruption dominated by MHD, E partitioning by kinetics
- Number of islands scales  $\sim S$
- For large  $S$  may dominate particle acceleration, as in Drake et al.

- Flare reconnection the central process of major solar eruptions
  - Little evidence for ideal processes in any of our work
  - Emphasizes key role of reconnection – similar to magnetosphere
- Need to validate/refute with observations
- Energy partitioning determined by reconnection dynamics
  - Will incorporate multiscale coupling into models
  - Anticipate major advances in both basic understanding and space weather modeling